

CONTENTS

(Full texts are incorporated in CJESEVIER, a file in the Chemical Journals Online database available on STN International; Abstracted, indexed in: Aluminium Abstracts; Anal. Abstr.; Biol. Abstr.; BIOSIS; Chem. Abstr.; Curr. Contents Phys. Chem. Earth Sci.; Engineered Materials Abstracts; Excerpta Medica; Index Med.; Life Sci.; Mass Spectrum. Bull.; Material Business Alerts; Metals Abstracts; Sci. Citation Index)

Special Issue


Papers presented at the 4th International Symposium on Hormone and Veterinary Drug Residue Analysis, Antwerp, Belgium, 4–7 June 2002. Part II

Determination of drug residues in feeds and animal products

- Rapid method for the determination of stanzolol in meat with supercritical fluid extraction and liquid chromatography–mass spectrometry
A.A.M. Stolker, E.F. van Tricht, P.W. Zoontjes, L.A. van Ginkel and R.W. Stephany (Bilthoven, The Netherlands) 1
- Value of alternative sample matrices in residue analysis for stanzolol
M.W.F. Nielen, H. Hooijerink, M.L. Essers, J.J.P. Lasaroms, E.O. van Bennekom and L. Brouwer (Wageningen, The Netherlands) 11
- Determination of propoxur residues in eggs by liquid chromatography–diode array detection after treatment of stocked housing facilities for the poultry red mite (*Dermanyssus gallinae*)
G. Hamscher, B. Prieß, J. Hartung, M.I. Nogosse, G. Glünder and H. Nau (Hannover, Germany) 19
- Simultaneous detection of five coccidiostats in eggs by liquid chromatography–tandem mass spectrometry
L. Mortier, E. Daeseleire (Melle, Belgium) and P. Delahaut (Marloie, Belgium) 27
- Multiresidue determination of fluoroquinolone antibiotics in eggs using liquid chromatography–fluorescence–mass spectrometryⁿ
M.J. Schneider (Wyndmoor, PA, USA) and D.J. Donoghue (Fayetteville, AR, USA) 39
- Screening for gestagens in kidney fat using accelerated solvent extraction and liquid chromatography electrospray tandem mass spectrometry
H. Hooijerink, E.O. van Bennekom and M.W.F. Nielen (Wageningen, The Netherlands) 51
- Screening procedures for clenbuterol residue determination in bovine urine and liver matrices using enzyme-linked immunosorbent assay and liquid chromatography
A. Posnyak, J. Zmudzki and J. Niedzielska (Pulawy, Poland) 61
- Liquid chromatography with fluorescence detection of amoxicillin and ampicillin in feeds using pre-column derivatization
V. Gamba and G. Dusi (Brescia, Italy) 69
- Food processing effects on residues: penicillins in milk and yoghurt
L. Grunwald and M. Petz (Wuppertal, Germany) 73
- Simultaneous determination of 13 quinolones from feeds using accelerated solvent extraction and liquid chromatography
I. Pecorelli, R. Galarini, R. Bibi, A. Floridi, E. Casciarri and A. Floridi (Perugia, Italy) 81
- Isolation of bound residues of nitrofurans from tissue by solid-phase extraction with determination by liquid chromatography with UV and tandem mass spectrometric detection
A. Conneely, A. Nugent, M. O'Keeffe (Dublin, Ireland), P.P.J. Mulder, J.A. van Rhijn (Wageningen, The Netherlands), L. Kovacsics, A. Fodor (Budapest, Hungary), R.J. McCracken and D.G. Kennedy (Belfast, UK) 91
- Liquid chromatographic–tandem mass spectrometric detection of banned antibacterial growth promoters in animal feed
C. Van Poucke, K. De Keyser (Ghent, Belgium), A. Baltusnikiene (Kaunas, Lithuania), J.D.G. McEvoy (Belfast, UK) and C. Van Peteghem (Ghent, Belgium) 99
- Liquid chromatographic–electrospray tandem mass spectrometric method for the determination of mebendazole and its hydrolysed and reduced metabolites in sheep muscle
H.D. Ruyck, E. Daeseleire, H. De Ridder and R. Van Renterghem (Melle, Belgium) 111

Determination of chloramphenicol in animal tissues and urine. Liquid chromatography–tandem mass spectrometry versus gas chromatography–mass spectrometry A. Gantverg, I. Shishani (Mississauga, Canada) and M. Hoffman (Washington, DC, USA)	125
Determination of ractopamine in animal tissues by liquid chromatography–fluorescence and liquid chromatography/tandem mass spectrometry E. Shishani, S.C. Chai, S. Jamokha, G. Aznar (Mississauga, Canada) and M.K. Hoffman (Washington, DC, USA)	137
Application of size-exclusion chromatography to the analysis of shrimp for sulfonamide residues J.E. Roybal, A.P. Pfenning, S.B. Turnipseed and S.A. Gonzales (Denver, CO, USA)	147
Immunoassays	
Screening and confirmation of chloramphenicol in shrimp tissue using ELISA in combination with GC–MS ² and LC–MS ² S. Impens (Merelbeke, Belgium), W. Reybroeck (Melle, Belgium), J. Vercammen, D. Courtheyn (Gentbrugge, Belgium), S. Ooghe (Melle, Belgium), K. De Wasch (Merelbeke, Belgium), W. Smedts (Brussel, Belgium) and H. De Brabander (Merelbeke, Belgium)	153
Surface plasmon resonance biosensor screening of poultry liver and eggs for nicarbazin residues B. McCarney, I.M. Traynor, T.L. Fodey, S.R.H. Crooks and C.T. Elliott (Belfast, UK)	165
Biosensor immunoassay for the detection of eight sulfonamides in chicken serum W. Haasnoot, M. Bienenmann-Ploum (Wageningen, The Netherlands) and F. Kohen (Rehovot, Israel)	171
Detection of levamisole residues in bovine liver and milk by immunobiosensor S.R.H. Crooks, B. McCarney, I.M. Traynor, C.S. Thompson, S. Floyd and C.T. Elliott (Belfast, UK)	181
Detection of multi- β -agonist residues in liver matrix by use of a surface plasma resonance biosensor I.M. Traynor, S.R.H. Crooks, J. Bowers and C.T. Elliott (Belfast, UK)	187
Production and characterisation of polyclonal antibodies to a range of nitroimidazoles T.L. Fodey, L. Connolly, S.R.H. Crooks (Belfast, UK), P. Delahaut (Marloie, Belgium) and C.T. Elliott (Belfast, UK)	193
Determination of drug residues in blood	
Analysis for endogenous and recombinant porcine somatotropine in serum M.H. Blokland, S.S. Sterk, L.A. van Ginkel, R.W. Stephany (Bilthoven, The Netherlands) and A.J.R. Heck (Utrecht, The Netherlands)	201
Clenbuterol food poisoning diagnosis by gas chromatography–mass spectrometric serum analysis F. Ramos (Coimbra, Portugal), A. Cristino (Lisbon, Portugal), P. Carrola, T. Eloy, J.M. Silva, M. da Conceição Castilho and M.I. Noronha da Silveira (Coimbra, Portugal)	207
Liquid chromatographic determination of levamisole in animal plasma: ultraviolet versus tandem mass spectrometric detection S. De Baere, M. Cherlet, S. Croubels, K. Baert and P. De Backer (Merelbeke, Belgium)	215
Exposure and withdrawal period	
Effect of long-term exposure to combinations of growth promoters in Long Evans rats. Part 1: Endocrine ovarian function J.C. Illera, G. Silván, M.M. Martínez-Mateos, A. Blass, P.L. Lorenzo and M. Illera (Madrid, Spain)	225
Effect of long-term exposure to combinations of growth promoters in Long Evans rats. Part 2: ovarian morphology J.C. Illera, G. Silván, M.M. Martínez-Mateos (Madrid, Spain), A.J. Conley, J. Corbin (Davis, CA, USA), A. Blass, P.L. Lorenzo and M. Illera (Madrid, Spain)	233
Influence of milking frequency on withdrawal period after application of β -lactam antibiotic-based drugs K. Knappstein, G. Suhren and H.-G. Walte (Kiel, Germany)	241
Estimation the withdrawal period for veterinary drugs used in food producing animals M.L. Vranic, L. Marangunic, H. Fernández Courel and A. Fernández Suárez (Buenos Aires, Argentina)	251
Steroids	
Application of neutral loss mode in liquid chromatography–mass spectrometry for the determination of corticosteroids in bovine urine E. Sangiorgi, M. Curatolo, W. Assini and E. Bozzoni (Brescia, Italy)	259
Faster analysis of anabolic steroids in kidney fat by downscaling the sample size and using gas chromatography–tandem mass spectrometry S. Impens (Merelbeke, Belgium), D. Courtheyn (Gentbrugge, Belgium), K.D. Wasch and H.F. De Brabander (Merelbeke, Belgium)	269
Oxidative stress as a bio-marker of estrogen exposure in healthy veal calves G. Brambilla, A. Ballerini, C. Civitareale, M. Fiori, B. Neri, R. Cavallina, A. Nardoni and L. Giannetti (Rome, Italy)	281
Modification of 17 β -estradiol metabolite profile in steer edible tissues after estradiol implant administration D. Maume, B. Le Bizec, K. Pouponneau, Y. Deceuninck, V. Solere (Nantes, France), A. Paris (Toulouse, France), J.-P. Antignac and F. Andre (Nantes, France)	289

Identification of anabolic steroids in serum, urine, sweat and hair. Comparison of metabolic patterns D. Thieme, P. Anielski, J. Grosse (Kreisch, Germany), H. Sachs (Munich, Germany) and R.K. Mueller (Kreisch, Germany)	299
Sample preparation	
Evaluation of sample preparation for control of chloramphenicol residues in porcine tissues by enzyme-linked immunosorbent assay and liquid chromatography A. Posyniak, J. Zmudzki and J. Niedzielska (Pulawy, Poland)	307
Development and optimisation of a method for the extraction of benzimidazoles from animal liver using supercritical carbon dioxide M. Danaher (Dublin, Ireland and Cork, Ireland), M. O'Keeffe (Dublin, Ireland) and J.D. Glennon (Cork, Ireland)	313
Validation	
Validation of analytical methods based on mass spectrometric detection according to the "2002/657/EC" European decision: guideline and application J.-P. Antignac, B. Le Bizec, F. Monteau and F. Andre (Nantes, France)	325
Validation of a method for detecting and quantifying tranquilisers and a β -blocker in pig tissues by liquid chromatography-tandem mass spectrometry Ph. Delahaut, C. Levaux, P. Eloy and M. Dubois (Marloie, Belgium)	335
Validation of a multiresidue liquid chromatography-tandem mass spectrometric method for the quantitation and confirmation of corticosteroid residues in urine, according to the proposed SANCO 1085 criteria for banned substances M.J. O'Keeffe, S. Martin and L. Regan (Dublin, Ireland)	341
Performance of a matrix-comprehensive in-house validation study by means of an especially designed software S. Uhlig, P. Gowik and W. Radeck (Berlin, Germany)	351
Screening, standards, calibration and stability	
Detection of cefquinome in milk by liquid chromatography and screening methods G. Suhren and K. Knappstein (Kiel, Germany)	363
Use of ion-trap liquid chromatography-mass spectrometry to screen and confirm drug residues in aquacultured products S.B. Turnipseed, J.E. Roybal, A.P. Pfenning (Denver, CO, USA) and P.J. Kijak (Laurel, MD, USA)	373
Identification of "unknown analytes" in injection sites: a semi-quantitative interpretation K. De Wasch, N. Van Hoof, S. Poelmans, L. Okerman (Merelbeke, Belgium), D. Courtheyn (Gentbrugge, Belgium), A. Ermens, M. Cornells (Brussel, Belgium) and H.F. De Brabander (Merelbeke, Belgium)	387
Production of clenbuterol, diethylstilbestrol and trenbolone mass standards in lyophilised bovine urine M.W. Pfaffi (Freising-Weihenstephan, Germany), B. Reck, R. Dreher (Darmstadt, Germany) and H.H.D. Meyer (Freising-Weihenstephan, Germany)	401
Estimation of the minimum detectable value for the determination of PCBs in fatty food samples by GC-ECD: a curvilinear calibration case J. Van Loco, V. Hanot, G. Huysmans, M. Elskens, J.M. Degroodt and H. Beernaert (Brussels, Belgium)	413
Practical approach for the stability testing of veterinary drugs in solutions and in biological matrices during storage S. Croubels, S. De Baere and P. De Backer (Merelbeke, Belgium)	419
Author Index	429
Calendar of forthcoming meetings	N1



<http://chemweb.com/gateways/elsevier.html>

CONTENTS
direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier Science books and journals. You can register for **ContentsDirect** online at: <http://contentsdirect.elsevier.com>



Full text of this journal is available, on-line from **ScienceDirect**. Visit www.sciencedirect.com for more information.
